

MUMFORD'S DEGREE OF CONTACT AND DIOPHANTINE APPROXIMATIONS

ROBERTO G. FERRETTI

INTRODUCTION

The Schmidt Subspace Theorem establishes that the solutions of some particular systems of diophantine approximations in projective spaces accumulates on a finite number of linear subspaces ([10], Theorem V.1.D'). One may state the following question: given a subvariety X of a projective space $\mathbb{P}^n$, does there exists a system of diophantine approximations on $\mathbb{P}^n$ whose solutions are Zariski dense in $\mathbb{P}$ but lie in finitely many proper subvarieties of X ? One can gain insight into this problem using a theorem of G. Faltings and G. Wüstholz ([2], Theorem 7.3, or [1] for a quantitative version). Their construction requires the hypothesis that the sum of some expected values (see (1.3)) has to be large. This sum turns out to be proportional to a degree of contact of a weighted flag of sections over the variety X (see (1.6)). The degree of contact was first introduced by D. Mumford (see for example [9]) as a tool for the construction of moduli spaces for families of algebraic schemes ([9] for curves, [4] for canonically embedded minimal surfaces, or [8] for ruled surfaces). Indeed, it measures the semistability of the Chow (or Hilbert) point of X under the action of an appropriate reductive algebraic group. Whence, the lower bound in the Faltings-Wüstholz theorem may be translated into a geometric invariant theoretic language. Moreover, the question of understanding when a system of diophantine approximations on X is not under control of the Schmidt Subspace Theorem relies on the Chow-unstability of X^s , for some sufficiently large $s > 0$. Unstability is usually associated to singular varieties. We will see this for some unstable local rings and for a family of elliptic surfaces. However ruled surfaces and some blow-ups give examples of non singular varieties having an unstable embedding. In general it is quite difficult to find examples of unstable projective varieties. All noteworthy instances we found have a common style: the destabilizing flags are constructed by the vanishing of sections along special divisors. Moreover, in order to control all the dimensions involved, it is crucial to use Riemann-Roch, as well as some result on the vanishing of higher cohomologies. It is a pleasure to thank J.-B. Bost, M. McQuillan, C. Soulé, G. Wüstholz for several discussions and remarks; IAS and IHES for the ospitality; and Swiss National Science Foundation and NSF with the grant DMS 9304580 for support.

1. PRELIMINARIES

Date: February 1, 2008.

1.1. Let K be an algebraic number field. Denote its ring of integers by $\mathcal{O}_K$ and its collection of places (equivalence classes of absolute values) by M_K . For $v \in M_K$, $x \in K$, we define the absolute value $|x|_v$ by

1. $|x|_v = |\sigma(x)|^{1/[K:\mathbb{Q}]}$ if v corresponds to the embedding $\sigma : K \hookrightarrow \mathbb{R}$;
2. $|x|_v = |\sigma(x)|^{2/[K:\mathbb{Q}]} = |\bar{\sigma}(x)|^{2/[K:\mathbb{Q}]}$ if v corresponds to the pair of conjugate embeddings $\sigma, \bar{\sigma} : K \hookrightarrow \mathbb{C}$;
3. $|x|_v = (N\mathfrak{p})^{-\text{ord}_{\mathfrak{p}}(x)/[K:\mathbb{Q}]}$ if v corresponds to the prime ideal $\mathfrak{p}$ of $\mathcal{O}_K$.

Here $N\mathfrak{p} = \#(\mathcal{O}_K/\mathfrak{p})$ is the norm of $\mathfrak{p}$ and $\text{ord}_{\mathfrak{p}}(x)$ the exponent of $\mathfrak{p}$ in the prime ideal decomposition of (x) , with $\text{ord}_{\mathfrak{p}}(0) := \infty$. In case (i) or (ii) we call v real infinite or complex infinite, respectively and write $v|\infty$; in case (iii) we call v finite and write $v \nmid \infty$. These absolute values satisfy the *Product Formula*

$$(1.1) \quad \prod_{v \in M_K} |x|_v = 1, \text{ for } x \in K^*.$$

The height of $\mathbf{x} = (x_0, \dots, x_N) \in K^{N+1}$ with $\mathbf{x} \neq 0$ is defined as follows: for $v \in M_K$ put

$$\begin{aligned} |\mathbf{x}|_v &= \left(\sum_{i=0}^N |x_i|_v^{2/[K:\mathbb{Q}]} \right)^{1/2[K:\mathbb{Q}]} \text{ if } v \text{ is real infinite,} \\ |\mathbf{x}|_v &= \left(\sum_{i=0}^N |x_i|_v^{[K:\mathbb{Q}]} \right)^{1/[K:\mathbb{Q}]} \text{ if } v \text{ is complex infinite,} \\ |\mathbf{x}|_v &= \max\{|x_0|_v, \dots, |x_N|_v\} \text{ if } v \text{ is finite.} \end{aligned}$$

Now define

$$h(\mathbf{x}) = h(x_0, \dots, x_N) = \sum_{v \in M_K} \log |\mathbf{x}|_v.$$

By the Product Formula (1.1) this defines a function on $\mathbb{P}^N(K)$. Further, $h(\mathbf{x})$ depends only on $\mathbf{x}$ and not on the choice of the number field K containing the coordinates of $\mathbf{x}$, in other words the function $h(\mathbf{x})$ extends to a function on $\mathbb{P}^N(\overline{\mathbb{Q}})$, where $\overline{\mathbb{Q}}$ is the algebraic closure of $\mathbb{Q}$ inside $\mathbb{C}$.

It is possible to define the height $h(X)$ of a projective variety $X \subseteq \mathbb{P}^N$ defined over K ([12] for a good reference). This invariant is comparable to the height of the coefficients of the (Cayley-Bertini-van der Waerden-)Chow point of X .

1.2. Let E be a vector space of rank $N + 1$ over the number field K and $E^\vee = \text{Hom}(E, K)$. Consider a closed subvariety $X \subseteq \mathbb{P}(E^\vee)$ in the projective space of lines of $E^\vee$, of dimension d . We choose a basis $l_0, \dots, l_N \in E$, fix non negative real numbers $r_0 \geq \dots \geq r_N$ and let $\mathbf{r} = (r_0, \dots, r_N)$. When m is large enough, $m \geq m_0$ say, the cup product map

$$\varphi : E^{\otimes m} \rightarrow H^0(X, \mathcal{O}(m))$$

is surjective, so that $H^0(X, \mathcal{O}(m))$ is generated by the monomials

$$l_0^{\alpha_0} \dots l_N^{\alpha_N} = \varphi(l_0^{\otimes \alpha_0} \otimes \dots \otimes l_N^{\otimes \alpha_N}),$$

with $\alpha_0 + \cdots + \alpha_N = m$. A special basis is a basis of $H^0(X, \mathcal{O}(m))$ made of such elements. We define the weight of l_i to be r_i , $i = 0, \dots, N$, the weight of a monomial in $E^{\otimes m}$ to be the sum of the weights of the l_i 's occurring in it, and the weight of a monomial $u \in H^0(X, \mathcal{O}(m))$ to be the minimum $w_{\mathbf{r}}(u)$ of the weights of the monomials in the l_i 's mapping to u by φ . The weight of a special basis is the sum of the weights of its elements. Finally, $w_{\mathbf{r}}(m)$ denotes the minimal weight among all special bases of $H^0(X, \mathcal{O}(m))$. Fix a special basis with minimal weight and denote by $w_1, \dots, w_M$ the weights of its elements in increasing order. Let F^j be the subspace of $H^0(X, \mathcal{O}(m))$, $j = 1, \dots, M$, generated by those monomials u for which the weight $w_{\mathbf{r}}(u)$ is smaller than w_j . We define a probability measure on $\mathbb{R}$ through the density function:

$$(1.2) \quad \rho_m(x) = \frac{1}{h^0(X, \mathcal{O}(m))} \sum_{w_j \leq x \cdot m} \dim(F^{j+1}/F^j) = \min_{w_j \leq x \cdot m} \frac{\dim F^j}{h^0(X, \mathcal{O}(m))}.$$

Then the expected value $E(\rho_m)$ of the probability measure is

$$(1.3) \quad E(\rho_m) = \frac{w_{\mathbf{r}}(m)}{m \cdot h^0(X, \mathcal{O}(m))}.$$

1.3. For integers $r_0 \geq \cdots \geq r_N$ and a basis $l_0, \dots, l_N$ of E , the *degree of contact* to X is the integer $e_{\mathbf{r}}(X)$, such that when m goes to infinity

$$(1.4) \quad w_{\mathbf{r}}(m) = e_{\mathbf{r}}(X) \frac{m^{d+1}}{(d+1)!} + O(m^d),$$

([8], Corollary 3.3). By the usual theory of Hilbert polynomials

$$(1.5) \quad h^0(X, \mathcal{O}(m)) = \deg(X) \frac{m^d}{d!} + O(m^{d-1}).$$

Combining (1.4) with (1.5) we obtain

$$(1.6) \quad E_{\infty}(X) := \lim_{m \rightarrow \infty} E(\rho_m) = \frac{e_{\mathbf{r}}(X)}{(d+1) \deg(X)}.$$

1.4. Let $L \supseteq K$ a finite field extension and Σ be a finite set of places of L containing all infinite places. For each $v \in \Sigma$ we choose a basis $l_{v,0}, \dots, l_{v,N}$ of $E \otimes_K L$ and non negative real numbers $r_{v,0} \geq \cdots \geq r_{v,N}$ as above. This defines, for all big enough integers m , a probability measure (1.2) and consequently a real number $E_{v,\infty}(X \otimes_K L)$ as in (1.6).

Theorem 1.5. *Let us suppose that for some real number $0 < \delta < 1$ we have,*

$$(1.7) \quad \sum_{s \in \Sigma} E_{s,\infty}(X \otimes_K L) = 1 + \delta.$$

Then there are two effectively computable constants c_1, c_2 depending only on $K, \Sigma, \delta, N, \deg(X), h(X)$, and the set $\{l_{v,i}; v \in \Sigma, i = 0, \dots, N\}$ such that all points $\mathbf{x} \in X(K)$ with

$$(1.8) \quad h(\mathbf{x}) > c_1,$$

and

$$(1.9) \quad \log\left(\frac{|l_{v,i}(\mathbf{x})|_v}{|\mathbf{x}|_v}\right) \leq -r_{v,i}h(\mathbf{x}) \quad v \in \Sigma, \quad i = 0, \dots, N,$$

are contained in at most c_2 proper subvarieties of X of degree not exceeding $\deg(X)$.

Proof. [1] Theorem 10.2 and [2] Theorem 7.3. $\square$

Remark 1.6. For this theorem it is allowed that all weights $r_{v,i}$ are rationals, but (1.7) still holds. This is no restriction since everything can be multiplied by a common denominator. We will tacitly assume this in the next sections.

1.7. There is an intersection theoretic formula expressing the degree of contact as the degree of divisor on a suitable modification of X .

Let $\mathcal{O}_{X \times \mathbb{A}^1}(1) = \mathcal{O}_X(1) \otimes \mathcal{O}_{\mathbb{A}^1}$ and t the coordinate of $\mathbb{A}^1$. For integers $r_0 \geq \dots \geq r_N$ and a basis $l_0, \dots, l_N$ of E we associate the $K[t]$ -submodule I of $H^0(X \times \mathbb{A}^1, \mathcal{O}_{X \times \mathbb{A}^1}(1))$ generated by $\{t^{r_i}l_i, \quad i = 0, \dots, N\}$ and an ideal sheaf $J \subseteq \mathcal{O}_{X \times \mathbb{A}^1}$ defined by

$$J \cdot \mathcal{O}_{X \times \mathbb{A}^1}(1) = \text{sheaf generated by } I \text{ in } \mathcal{O}_{X \times \mathbb{A}^1}(1).$$

Choose a compactification Y of $X \times \mathbb{A}^1$ to which $\mathcal{O}_{X \times \mathbb{A}^1}(1)$ extends to a line bundle $\mathcal{L}$ and let $\pi : B \rightarrow Y$ be a blow-up for which $\pi^{-1}(J) = \mathcal{O}_B(-E)$, where E is the exceptional divisor. Let $\pi^{-1}(\mathcal{L}) = \mathcal{O}_B(D)$, then we have

$$(1.10) \quad e_{\mathbf{r}}(X) = D^d - (D - E)^d,$$

([8], §2; [9], §2). One checks that the property is independent of the choices involved. Let Z be the subscheme of $X \times \mathbb{A}^1$ defined by the ideal sheaf J . According [3] §4, one may write the degree of contact in terms of Segre classes:

$$e_{\mathbf{r}}(X) = \sum_{k=1}^d (-1)^k \binom{d}{k} L^{d-k} s_k(Z, Y),$$

([3], Corollary 4.2.2 and projection formula). Thus, when Z is set-theoretically a point $w_{\mathbf{r}}(m)$ is the Hilbert-Samuel polynomial of J as an ideal of $\mathcal{O}_{X \times \mathbb{A}^1}$ and $e_{\mathbf{r}}(X)$ the multiplicity there ([9], §2 Examples).

1.8. The Chow point of X is *semistable* with respect to integers $r_0 \geq \dots \geq r_N$ and a basis $l_0, \dots, l_N$ of E if and only if

$$(1.11) \quad E_{\infty}(X) \leq \frac{1}{n+1} \sum_{i=1}^N r_i$$

([9], Theorem 2.9). If this properties is satisfied for all bases and weights as above, the Hilbert-Mumford theorem implies that the Chow point of X is semistable with respect to the action of $SL(E)$. It is needless to say that the opposite of semistable is *unstable*.

1.9. We finish this section with an important remark. Schmidt Subspace Theorem ([10], Theorem V.1.D') corresponds to Theorem 1.5 when X is the full projective space $\mathbb{P}^N$. We would like that Theorem 1.5 for an arbitrary projective variety $X \subseteq \mathbb{P}^N$ is not a simple corollary of Schmidt Subspace Theorem for $\mathbb{P}^N$. Paragraph **1.8** may help us with this task. It shows that the possible varieties where this can be avoided are those for which the Chow point of $X \times \cdots \times X$ ($\# \Sigma$ times) is unstable.

2. LOCAL RINGS

2.1. Let $K \subseteq L$ be number fields, Σ a finite set of places of L containing all infinite places. Let E be a vector space of rank $N+1$ over K and $E^\vee = \text{Hom}(E, K)$. Consider a closed subvariety $X \subseteq \mathbb{P}(E^\vee)$ in the projective space of lines of $E^\vee$, of dimension of d . Let $P \in X(L)$ be a closed point, and consider a basis $l_0, \dots, l_N$ of $E \otimes_K L$ so that $P = (1, 0, \dots, 0)$ with respect to the coordinates in $\mathbb{P}(E^\vee)$ defined by this basis. For $v \in \Sigma$ we define weights $r_{0,v} = k_v$, $r_{1,v} = \dots = r_{N,v} = 0$.

Proposition 2.2. *Let $P \in X(L)$ be a closed point and for $v \in \Sigma$ let $d_v > 0$ real such that*

$$\text{mult}_P X \sum_{v \in \Sigma} \frac{k_v}{d_v} > (\dim X + 1) \deg X.$$

Then the points $\mathbf{x} \in X(K)$ with

$$\log \frac{|l_0(\mathbf{x})|_v}{|\mathbf{x}|_v} \leq -\frac{k_v}{s_v} h(\mathbf{x}), \quad v \in \Sigma,$$

lie in finitely many subvarieties of X of bounded degree

Proof. For each $v \in \Sigma$ the ideal $J \cdot \mathcal{O}_{X \times \mathbb{A}^1}(1)$ of Proposition 1.7 is generated by $\{t^{k_v} l_0, l_1, \dots, l_N\}$. Since $\{l_1, \dots, l_N\}$ generate the maximal ideal $\mathfrak{m}_{P,X}$ and l_0 is a unit at P , $J = (t^{k_v}, \mathfrak{m}_{P,X}) \mathcal{O}_{X \times \mathbb{A}^1}(1)$. Hence

$$e_{\mathbf{r}}(X) = k_v \cdot \text{mult}_{(0,P)}(X \times \mathbb{A}^1) = k_v \cdot \text{mult}_P X.$$

The claim follows by the linearity of the expected value in the weights, (1.6) and Theorem 1.5. $\square$

2.3. The attempt to generalize this theorem leads to a numerical measure of the degree of singularity of a point. Let R be a local ring of dimension r and m a positive integer, then the m -th flat multiplicity $e_m(R)$ of R is defined by

$$\begin{aligned} e_0(R) &= \sup \left\{ \frac{e(I)}{r! \text{col}(I)} : I \text{ of finite colength in } R \right\} \\ e_m(R) &= e_0(R[[t_0, \dots, t_m]]), \end{aligned}$$

where $e(I)$ denotes the multiplicity of the ideal I . Further, a local ring is called *semistable* if $e_1(R) = 1$. Let X be a projective scheme. Let L be an ample line bundle on X . For a sufficiently large n let

$$\phi_n : X \rightarrow \mathbb{P}^{h^0(X, L^{\otimes n}) - 1},$$

be the embedding defined by $L^{\otimes n}$. Suppose that there exists a point P on X such that the local ring $\mathcal{O}_{P,X}$ is unstable. Then for every positive integer n , there exists a positive integer $m > n$ such that the Chow point and the Hilbert point corresponding to $\phi_n(X)$ are unstable under the natural action of $SL(h^0(X, L^{\otimes m}))$ ([9], Proposition 3.12, [11], Proposition 1.3). Little is known on the semistability of local rings. See however [11] and [9], §3.

2.4. Let $K \subseteq L$ be number fields and Σ a finite set of places of L containing all infinite places. Let X be a projective scheme defined over K and $\mathcal{L}$ an ample line bundle on X . Suppose that there is a point closed P on $X(L)$ such that the local ring $\mathcal{O}_{P,X}$ is unstable. By [9], Lemma 3.6 there exists a sequence of ideals of finite colength

$$I_0 \subseteq I_1 \subseteq \cdots \subseteq I_N = \mathcal{O}_{P,X} = I_{N+1} = \cdots$$

such that if I is the ideal $\bigoplus_{i \geq 0} I_i t^i \subseteq \mathcal{O}_{P,X}[[t]]$, then

$$(2.1) \quad e(I) = (1 + \varepsilon)(\dim X + 1)! \operatorname{col}(I),$$

where $\varepsilon > 0$. By Riemann-Roch on X we can choose n sufficiently large so that

1. $\mathcal{L}^{\otimes n}$ is very ample,
2. the map $S^m H^0(X, \mathcal{L}^{\otimes n}) \rightarrow H^0(X, \mathcal{L}^{\otimes nm})$ is surjective for all $m \geq 1$,
3. the map $\phi_m : H^0(X, \mathcal{L}^{\otimes mn}) \rightarrow \mathcal{O}_{P,X}/I_0^m$ is surjective for all $m \geq 1$.

The vector space $H^0(X, \mathcal{L}^{\otimes n})$ has the induced filtration defined by

$$F^j = \phi_i^{-1}(I_j/I_0), \quad i = 0, \dots, N.$$

For each $v \in \Sigma$ choose a basis $l_{v,0}, \dots, l_{v,N}$ of $H^0(X, \mathcal{L}^{\otimes n})$ compatible with the filtration. Assign weights as in (3.2),

$$(2.2) \quad r_{v,i} = \min\{j : l_{v,i} \in F^j\}.$$

Theorem 2.5. *For $v \in \Sigma$ let $d_v > 0$ be reals such that*

$$(2.3) \quad \sum_{v \in \Sigma} \sum_{i=0}^N \frac{r_{v,i}}{s_v} > \frac{\deg X}{(\dim X + 1)!}.$$

Then the points $\mathbf{x} \in X(K)$ with

$$\log\left(\frac{|l_{v,i}(\mathbf{x})|_v}{|\mathbf{x}|_v}\right) \leq -\frac{r_{v,i}}{d_v} h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, \dots, N$$

are contained in finitely many subvarieties of X of bounded degree.

Proof. Since the colength of I corresponds to the sum of weights ([11], p. 334), the theorem follows by (2.1), (2.3) and Theorem 1.5. $\square$

Remark 2.6. According to [11] Appendix A, and 1.9, Theorem 2.5 is stronger than the Schmidt Subspace Theorem. For several examples of destabilizing weighted flags of unstable two-dimensional local rings, we refer to [11], §§4, 5, 6.

3. RULED SURFACES

3.1. The Steiner surface $X \subseteq \mathbb{P}^4$ is the closure of the image of the map $\psi : \mathbb{P}^2 \rightarrow \mathbb{P}^4$ defined by

$$(x, y, z) \mapsto (xz, yz, x^2, xy, y^2) = (l_0, \dots, l_4).$$

This surface is the blown up at the point $P = (0, 0, 1)$ and embedded by the system of conics passing through P . Its degree is three, the number of free points of intersection of two such conics, and it is a rational ruled surface of type $\mathbb{F}_1$, ruled by the pencil of lines passing through P . Let us consider the flag given by choosing the linear forms l_i as above and with weights $r_0 = r_1 = k$ for some $k > 0$ and $r_2 = r_3 = r_4 = 0$. Then then a computation shows that $e_r(X) = 4k$ ([8], Example 3.6). Theorem 1.5 implies:

Proposition 3.2. *Let $K \subseteq L$ be number fields, Σ a finite set of places of L containing all infinite places and for each $v \in \Sigma$ non-negative integers k_v and positive rationals d_v with*

$$\sum_{v \in \Sigma} \frac{k_v}{d_v} > \frac{9}{4}.$$

The solutions $\mathbf{x} \in X(K)$ of the inequalities

$$\log\left(\frac{|l_i(\mathbf{x})|_v}{|\mathbf{x}|_v}\right) \leq -\frac{k_v}{d_v}h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, 1$$

are contained in finitely many curves of X .

Remark 3.3. If we consider the same problem for $\mathbf{x} \in \mathbb{P}^4(K)$, then Schmidt Subspace Theorem implies that its solutions are contained in finitely many subspaces of $\mathbb{P}^4$, hence in finitely many curves of X , if the condition $\sum_{v \in S} (k_v/d_v) > 5/2 (> 9/4)$ holds true.

3.4. A good reference for the general properties of ruled surfaces is [5], V.2. Fix a smooth curve C of genus g and a geometrically ruled surface $p : R \rightarrow C$. For one section $\sigma : C \rightarrow R$ of p we refer to the divisor S on R which is the image of σ . Fix one section, say S , and let f denote the numerical equivalence class of a fibre of p . When the parity of S^2 , which is independent of S , is even. A convenient basis of $\text{Num}(R)$ is given by f and the element G determined by $G^2 = 0$ and $G \cdot f = 1$. For surfaces of odd parity such a G can be found in $\text{Num}(R) \otimes \mathbb{Q}$. We will by abuse of language consider G as a divisor on R . This is no restriction since we will be utilizing only the numerical properties of G .

3.5. Fix a ruled surface R , a very ample divisor $D \sim aG + bf$ on R , and a section S such that $D - aS$ is effective. Let $\mathbb{P}^N = \mathbb{P}(H^0(R, D)^\vee)$ and let $F^j = \{ h \in H^0(R, D), h \in H^0(R, D - jS) \}$. Since a section in D can vanish to order at most a on S ,

$$(3.1) \quad H^0(R, D) = F^0 \supset F^1 \supset \dots \supset F^a \supset F^{a+1} = \{0\}.$$

Choose a basis $l_0, \dots, l_N$ of $H^0(R, D)$ compatible with this flag, and fix weights r_i , $i = 0, \dots, N$ by the condition that

$$(3.2) \quad r_i = a - j \iff x_i \in F^j \setminus F^{j+1}.$$

That is, r_i equals a minus the order to which l_i vanishes along S , for $i = 0, \dots, N$. Clearly, the r_i 's decrease to zero. This construction generalize **3.1** of the Steiner surface in $\mathbb{P}^4$. In that construction we have $C = \mathbb{P}^1$, $E = \mathcal{O}_{\mathbb{P}^1}(1) \otimes \mathcal{O}_{\mathbb{P}^1}(2)$ and $R = \mathbb{P}(E)$. Further, $D = \mathcal{O}_{\mathbb{P}(E)}(1) \sim G + 3/2f$ and S , the exceptional divisor, which is numerical equivalent to $G - 1/2f$ and which is the section associated to the bundle $\mathcal{O}_{\mathbb{P}^1}(2)$ in E , we get the flag of that example. By [8] Theorem 6.5 we obtain a non-trivial generalisation of Schmidt Subspace Theorem for this embedding:

Theorem 3.6. *Let $K \subseteq L$ be number fields and Σ a finite set of places of L containing all infinite places. Let S be a section of R . For each $v \in \Sigma$ let $l_{v,0}, \dots, l_{v,N}$ a basis compatible with the filtration (3.1) associated to S and let $r_{v,0}, \dots, r_{v,N}$ be the corresponding weights (3.2). Suppose that $H^i(R, D - jS) = 0$ for $i > 0$ and $0 \leq j \leq a$. Further, for each $v \in \Sigma$ let $d_v > 0$ reals such that*

$$(3.3) \quad (3a^2D \cdot S - a^3S^2) \sum_{v \in \Sigma} \frac{1}{d_v} > 3D^2.$$

Then the points $\mathbf{x} \in R(K)$ with

$$\log \frac{|l_i(\mathbf{x})|_v}{|\mathbf{x}|_v} \leq -\frac{r_{v,i}}{d_v} h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, \dots, N$$

lie in finitely many curves of R of bounded degree.

Proof. This follows by Theorem 1.5 and [8] Proposition 6.2. $\square$

3.7. If $p : E \rightarrow C$ is a rank 2 vector bundle on C such that $\mathbb{P}(E) \cong R$ we say that E represent R . Such E always exists and R determines E up to tensoring with a line bundle L on C . Let L be a line subbundle of E of maximal degree. The number $\deg E - 2 \deg L$ is independent of the choice of E . Then we say that the ruled surface is *bundle semistable* if $\deg E \geq 2 \deg L$ and *bundle unstable* when $\deg E < 2 \deg L$. In the latter case there exists a unique section S of negative selfintersection.

Theorem 3.8. *Let $K \subseteq L$ be number fields, Σ a finite set of places of L containing all infinite places. Suppose R is bundle unstable. Let $D \sim aG + bf$ be a very ample on R . Let S be the unique section of R of negative selfintersection. For $v \in \Sigma$ let $l_{v,0}, \dots, l_{v,N}$ a basis compatible with the filtration (3.1) associated to S and let $r_{v,0}, \dots, r_{v,N}$ be the corresponding weights (3.2). Further, for each $v \in \Sigma$ let $d_v > 0$ reals such that*

$$(3.4) \quad b + \frac{a}{2}(\deg E - 2 \deg L) > 2g - 2,$$

and

$$(3.5) \quad \sum_{v \in \Sigma} \frac{1}{d_v} > \frac{3b}{a(b + g - 1)}.$$

Then the points with $\mathbf{x} \in R(K)$ with

$$\log \frac{|l_i(\mathbf{x})|_v}{|\mathbf{x}|_v} \leq -\frac{r_{v,i}}{d_v} h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, \dots, N$$

lie in finitely many curves of R of bounded degree.

Proof. Since $\deg E < 2 \deg L$ and b is bounded below by (3.4), $H^i(R, D - jS) = 0$ for $i > 0$ and $0 \leq j \leq a$. ([8], Remark after Proposition 6.2). The degree of R is $D^2 = 2ab$. According to [8] (6.7) we have $3a^2D \cdot S - a^3S^2 = \frac{a^3}{2}l_R + 3a^2b$. The inequalities (3.4) and (3.5) imply (3.3) and Theorem 3.6 concludes the theorem. $\square$

3.9. Little is known of the unstability of higher dimensionals varieties. Each new information on this problem has always interesting consequences. For instance, Vojta's Conjectures on the ruled threefold defined by the equation $a^4x + b^4y + c^4z = 0$ in $\mathbb{P}^2 \times \mathbb{P}^2$ imply *abc*-conjecture.

4. BLOW-UP

4.1. Let $K \subseteq L$ be number fields, Σ a finite set of places of L containing all infinite places. Let X be a normal variety defined over K of dimension n , H a Cartier divisor on X , E an effctive Cartier divisor and fix an integer $s > 0$. Assume that for some sufficiently large r , larger than $r_0 \geq s$ say, $rH - sE$ is very ample. Put $W_{i,j} = H^0(X \otimes_K L, iH - jE)$ for $i, j \in \mathbb{Z}$. Then $W_{r,s}$ has a filtration $W_{r,s} \supset W_{r,s+1} \supset \cdots \supset W_{r,r}$. For $v \in \Sigma$ let $l_{v,0}, l_{v,1}, \dots, l_{v,N}$ be a basis of $W_{r,s}$ compatible with the filtration. For these bases we consider weights $r_{v,i} = r - j$ if $l_{v,i} \in W_{r,j}$ and $l_{v,i} \notin W_{r,j+1}$, $v \in \Sigma$.

Theorem 4.2. *Let us suppose that for any $i > 0$*

$$h^i(X \otimes_K L, rH - jE) = O((r+j)^{n-2}) \text{ on } \{(r,j) \in \mathbb{Z}^2 \mid s \leq j \leq r\}.$$

For $v \in \Sigma$ let d_v be positive reals with

$$\sum_{i=0}^n H^{n-i} \left(\frac{1}{4E} \right)^i \binom{n+1}{i+1} ((i+1)(r-s)r^{n-i}s^i - r^{n+1} + r^{n-i}s^{i+1}) \sum_{v \in \Sigma} \frac{1}{d_v} > (n+1)(rH - sE)^n. \quad (4.2)$$

Then the points $\mathbf{x} \in X(K)$ with

$$\log\left(\frac{|l_{v,i}(\mathbf{x})|_v}{|\mathbf{x}|_v}\right) \leq -\frac{r_{v,i}}{d_v}h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, \dots, N$$

are contained in finitely many subvarieties of X of bounded degree.

Proof. For each $v \in \Sigma$ the filtrations constructed above correspond to the filtration of [6], Lemma 3. In *loc. cit.* it is proven that, for each $v \in \Sigma$ the left hand side of (4.2) is a lower bound for the degree of contact. The assertion we want follows then by Theorem 1.5. $\square$

Remark 4.3. Lemma 3 of [6] confirms that this theorem is stronger than Schmidt Subspace Theorem restricted to X . Further, one should observe that all intersection products of (4.2) can be written down by Segre classes.

4.4. In this section we follow [6], §2. Let V be a normal projective variety of dimension $n - 1 \geq 1$ over K , D an ample divisor on V and $Y_t \hookrightarrow \mathbb{P}^{N(t)+1}$ the projective cone over $\Phi_{|tD|} : V \hookrightarrow \mathbb{P}^{N(t)}$. Let $f_t : X_t \rightarrow Y_t$ be the blow-up with center the vertex. Put $\mathcal{O}_{Y_t}(H_{0,t}) = \mathcal{O}_{\mathbb{P}^{N(t)+1}}(1)|_{Y_t}$, $H_t = f_t^*H_{0,t}$ and E_t the exceptional divisor of f_t .

Proposition 4.5. *Take a large integer t in such a way that*

$$(4.3) \quad H^i(V \otimes_K L, mt\mathcal{L}) = 0, \text{ for any } m > 0 \text{ and } i > 0.$$

Further, assume that

$$(4.4) \quad (n+1)s^n(s-r) + r^{n+1} - s^{n+1} > \frac{n+1}{\#\Sigma}(r^n - s^n).$$

Then the result of Theorem 4.2 holds.

Proof. Condition (4.3) implies the vanishing of higher cohomologies for each $rH_t - jE_t$, $j = 1, \dots, r$ ([6], Proposition 4). Moreover, all intersection numbers $H_t^{n-i}(-E_t)^i$ are zero except for $i = 0, n$ where $H_t^n = -(-E_t)^n$. Thus, (4.4) implies (4.2) and the proposition follows. $\square$

Remark 4.6. The projective cone over a non-singular conic in $\mathbb{P}^2$ and a non-singular quadratic surface in $\mathbb{P}^3$ satisfy the condition of Proposition 4.5 for each $t \geq 1$ ([6], Example 6.)

4.7. We consider here resolutions of singularities. Let $f : X \rightarrow Y$ be the blow-up of a normal projective variety Y whose center is a subscheme Z not necessarily irreducible or reduced. Assume X is non-singular of dimension n . Let E be the exceptional divisor. Put $H = f^*H_0$ for an ample divisor H_0 on Y , and fix an integer $s \geq 0$.

Proposition 4.8. *Take a sufficiently large integer t such that $tH - E$ is ample on X and tH_0 is very ample on Y . Assume (4.2), then the assertion of Theorem 4.2 holds.*

Proof. The vanishing for the higher cohomology spaces is assured by the proof of [6], Proposition 8. $\square$

5. ELLIPTIC SURFACES

5.1. Following [7], we call a flat proper map of $\overline{\mathbb{Q}}$ -schemes $p : X \rightarrow \mathbb{P}^1$ a rational Weierstrass fibration if X is reduced and irreducible rational surface over $\overline{\mathbb{Q}}$, every geometric fibre of p is an irreducible curve of genus 1, and a section $s : \mathbb{P}^1 \rightarrow X$ is given, not passing through the nodes or cusps of the fibres. Moreover, we will assume that X is normal, and that the generic fibre of p is smooth. In this case, we may resolve the singularities of X and obtain an elliptic surface $\overline{p} : \overline{X} \rightarrow \mathbb{P}^1$ (with section) which we call the induced elliptic surface. In this situation one may represent X in Weierstrass form by the equation

$$y^2 = x^3 + A(t)x + B(t),$$

where A is a quartic and B is a sextic polynomial in the parameter t of $\mathbb{P}^1$.

5.2. Let us fix the following situation: $\overline{X}$ is a minimal rational elliptic surface with section and a fibre of type IV^* (in Kodaira classification), X the associated Weierstrass fibration, S the given section of $\overline{X}$, f the numerical class of a fibre of $\overline{X}$, E the unique rational component of the singular IV^* fibre having multiplicity 3, $D = 3kS +$

$6kf$ a very ample divisor on X for some positive integer k . We have a filtration on $H^0(\overline{X}, \mathcal{O}(D))$ defined by

$$(5.1) \quad H^0(\overline{X}, \mathcal{O}(D)) \supset H^0(\overline{X}, \mathcal{O}(D - E)) \supset \cdots \supset H^0(\overline{X}, \mathcal{O}(D - 18kE)) \supset \{0\}.$$

Let K be a number field defining $\overline{X}$ as a geometrically irreducible variety, and L a finite extension. Let Σ a finite set of places of L containing all infinite places. For each $v \in \Sigma$, we choose a basis $l_{v,0}, \dots, l_{v,n}$ compatible with this filtration, and define weights $r_{v,i} = 18k - j$ if $l_{v,i}$ is in $H^0(\overline{X}, \mathcal{O}(D - jE))$ but not in $H^0(\overline{X}, \mathcal{O}(D - (j+1)E))$, $i = 0, \dots, n$.

Theorem 5.3. *For $v \in \Sigma$ let d_v be positive reals such that*

$$\sum_{v \in \Sigma} \frac{k}{d_v} > \frac{3}{25}.$$

Then all $\mathbf{x} \in X(K)$ with

$$\log\left(\frac{|l_{v,i}(\mathbf{x})|_v}{|\mathbf{x}|_v}\right) \leq -\frac{r_{v,i}}{d_v}h(\mathbf{x}), \quad v \in \Sigma, \quad i = 0, \dots, n$$

are contained in finitely many curves of X of bounded degree.

Proof. According to [7], (6.7) we know that the degree of contact of the filtration (5.1) is bounded below by $675k^3$. On the other hand, the degree of X with respect to D equals $27k^2$ ([7], (4.1)). Hence,

$$\frac{e_{\mathbf{r}}(X)}{(\dim X + 1) \deg X} \geq \frac{675k^3}{3 \cdot 27k^2} = \frac{25}{3}k.$$

Theorem 1.5 confirms the the theorem. $\square$

Remark 5.4. Since D is very ample, by Riemann-Roch we can compute the dimension $n + 1$ of the space $H^0(\overline{X} \otimes_K L, \mathcal{O}(D))$ which equals $(27k^2 + 3k + 2)/2$ ([7], (4.2)). For each $v \in \Sigma$

$$\sum_{i=0}^n r_{v,i} = \frac{225}{2}k^3 + 9k^2 + \frac{23}{2}k,$$

([7], (5.2)). Moreover, for all $k \geq 1$ we have

$$\frac{25}{3}k > \frac{1}{\frac{27k^2+3k+2}{2}}\left(\frac{225}{2}k^3 + 9k^2 + \frac{23}{2}k\right).$$

This means that Schmidt Subspace Theorem restricted to X is weaker than Theorem 5.3

Remark 5.5. The filtration (5.1) is clearly given by the order of vanishing for sections of D along the curve E . Calculations of the same type can be carried through for Weierstrass fibrations with singularities of type II^* and III^* . The filtrations used in these latter cases are also given by the order of vanishing along the unique curve of

maximum multiplicity in the singular fibre in question. This curve is the multiplicity six component in the III^* fibre and the multiplicity four component in the II^* fibre.

REFERENCES

- [1] R. G. FERRETTI, *Diophantine Approximations on Projective Varieties*, Preprint.
- [2] G. FALTINGS, G. WÜSTHOLZ, Diophantine Approximations on Projective Spaces, *Invent. Math.*, **116** (1994) ,109—138.
- [3] W. FULTON, *Intersection Theory*, Springer, 1984.
- [4] D. GIESEKER, Global moduli for surfaces of general type, *Invent. Math.*, **43** (1977) 233—282.
- [5] R. HARTSHORNE, *Algebraic Geometry*, Springer, 1977.
- [6] S. ISHII, Chow instability of certain projective varieties, *Nagoya Math. J.*, **92** (1983) 39—50.
- [7] R. MIRANDA, Projectively Unstable Elliptic Surfaces, *Illinois J. Math.*, **27** (1983) 404—420.
- [8] I. MORRISON, Projective Stability of Ruled Surfaces, *Invent. Math.*, **56** (1980) 269—304.
- [9] D. MUMFORD, Stability of Projective Varieties, *Enseign. Math.*, **XXIII** (1977) 39—110.
- [10] W. M. SCHMIDT, *Diophantine Approximations and Diophantine Equations*, LNM 1467, Springer 1991.
- [11] J. SHAH, Stability of Two-Dimensional Local Ring.I, *Invent. Math.*, **64** (1981) 297—343.
- [12] C. SOULÉ, ET. AL., *Lectures on Arakelov Geometry*, Cambridge University Press, 1992.

INSITUTE FOR ADVANCED STUDY, SCHOOL OF MATHEMATICS, OLDEN LANE, PRINCETON, NJ 08540

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, 35 ROUTE DE CHARTRES, 91440 BURES-SUR-YVETTE, FRANCE

E-mail address: ferretti@ias.edu, ferretti@ihes.fr